



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Foreign and insular statistical reports of countries and cities—Yearly and monthly.

AFRICA—*Sierra Leone*.—Two weeks ended April 5, 1901. Estimated population, 35,000. Total number of deaths, 22. No infectious or contagious diseases.

ARGENTINA—*Buenos Ayres*.—Month of February, 1901. Estimated population, 825,858. Total number of deaths, 1,058, including diphtheria, 9; enteric fever, 19; measles, 10; scarlet fever, 29; smallpox, 32, and 35 from tuberculosis.

BRAZIL—*Ceara*.—Month of February, 1901. Estimated population, 50,000. Total number of deaths, 165, including enteric fever, 12, and 2 from measles.

Month of March, 1901. Total number of deaths, 215, including enteric fever, 32, and 3 from measles.

BRITISH WEST INDIES—*Trinidad—Port of Spain*.—Four weeks ended March 30, 1901. Estimated population, 34,037. Total number of deaths, 109, including measles, 1; whooping cough, 2, and 14 from tuberculosis.

CHILE—*Antofagasta*.—Month of March, 1901. Estimated population, 16,000. Total number of deaths, 66, including whooping cough, 3, and 6 from phthisis pulmonalis.

FRANCE—*Nice*.—Month of March, 1901. Estimated population, 108,227. Total number of deaths, 257, including diphtheria, 4; enteric fever, 2; measles, 1; scarlet fever, 2, and 30 from tuberculosis.

St. Etienne.—Two weeks ended April 15, 1901. Estimated population, 146,030. Total number of deaths, 142, including diphtheria, 3, and 17 from tuberculosis.

GERMANY—*Dresden*.—Month of February, 1901. Estimated population, 398,650. Total number of deaths, 525, including diphtheria, 5; measles, 3; whooping cough, 6, and 67 from phthisis pulmonalis.

Kehl.—Month of March, 1901. Estimated population, 152,826. Total number of deaths, 301, including diphtheria, 6; whooping cough, 2; typhus fever, 1, and 39 from tuberculosis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended April 13, 1901, correspond to an annual rate of 19.2 per 1,000 of the aggregate population, which is estimated at 11,789,099. The highest rate was recorded in Liverpool, viz, 25.0, and the lowest in Croydon, viz, 11.2.

London.—One thousand six hundred and thirty-three deaths were registered during the week, including measles, 52; scarlet fever, 8; diphtheria, 26; whooping cough, 74; enteric fever, 4, and diarrhea and dysentery, 9. The deaths from all causes correspond to an annual rate of 18.3 per 1,000. In Greater London 2,162 deaths were registered, corresponding to an annual rate of 16.6 per 1,000 of the population.

In the "outer ring" the deaths included 8 from diphtheria, 14 from measles, 2 from scarlet fever, and 21 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended April 13, 1901, in the 22 principal town districts of Ireland was 27.9 per 1,000 of the population, which is estimated at 1,058,722. The lowest rate was recorded in Clonmel, viz, 9.7, and the highest in Tralee, viz, 61.6 per 1,000. In Dublin and suburbs 242 deaths were registered, including enteric fever, 2; scarlet fever, 2, and 8 from whooping cough.

Scotland.—The deaths registered in 8 principal towns during the week ended April 13, 1901, correspond to an annual rate of 23.2 per 1,000 of the population, which is estimated at 1,630,155. The lowest mortality was recorded in Perth, viz, 16.8, and the highest in Glasgow, viz, 24.1 per 1,000. The aggregate number of deaths registered from all causes was 726, including diphtheria, 4; measles, 17; scarlet fever, 3; smallpox, 6, and 51 from whooping cough.

JAPAN—Nagasaki.—Ten days ended March 31, 1901. Estimated population, 131,701. Number of deaths not reported. One death from enteric fever reported. No infectious or contagious disease prevailing.

MALTA.—Week ended April 6, 1901. Estimated population, 181,698. Total number of deaths, 76, including enteric fever, 1, and 2 from whooping cough.

Two weeks ended April 13, 1901. Estimated population, 181,698. Total number of deaths, 89, including diphtheria, 1, and 1 from whooping cough.

SPAIN—Malaga.—Two weeks ended March 31, 1901. Estimated population, 126,000. Total number of deaths, 198, including enteric fever, 1; leprosy, 1, and 3 from smallpox.

SWITZERLAND.—Reports from 18 cities and towns, for the week ended April 6, 1901, show a total of 310 deaths, including diphtheria, 1; enteric fever, 3; measles, 4; scarlet fever, 1; whooping cough, 2, and 33 from phthisis pulmonalis.